



Presentation Overview

Characteristics of Zika virus
 Characteristics of Zika virus infection
 Epidemiology and outbreaks
 Vector and transmission
 Clinical symptoms
 Diagnosis and testing
 Treatment and prevention
 Wisconsin response and preparedness
 Disease reporting and investigation
 Research at UW-Madison

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

2

Some Mosquito Borne Diseases

- | | |
|-------------------------------|----------------------------------|
| ▪ Chikungunya | ▪ Malaria |
| ▪ Dengue | ▪ Rift Valley fever |
| ▪ Eastern equine encephalitis | ▪ St. Louis encephalitis |
| ▪ Filariasis | ▪ Venezuelan equine encephalitis |
| ▪ Jamestown Canyon virus | ▪ Western equine encephalitis |
| ▪ Powassan virus | ▪ Yellow fever |
| ▪ Japanese encephalitis | ▪ ZIKA |

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

3

Zika Virus



- First identified in Uganda in 1947
- From the blood of a sentinel rhesus monkey during a study of sylvatic cycle of yellow fever virus
 - Febrile illness
- Isolated from mosquitoes in 1948
 - *Aedes africanus*

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

4

Zika Virus

- Mosquito-borne single-stranded RNA flavivirus
 - 10,794 bases long
 - Closely related to dengue, yellow fever, Japanese encephalitis and West Nile viruses.
 - 40 nm diameter
 - Icosahedral
 - Lipid enveloped
 - Neurotropic



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

5

Zika Virus Epidemiology

- Prior to 2007, only sporadic human cases reported from Africa and southeast Asia
 - Many cases and outbreaks likely not recognized
- 2007, first outbreak reported on Yap Island, Federated States of Micronesia
- 2013-2014, >28,000 suspected cases reported from French Polynesia
- May 2015, locally-acquired cases reported in Brazil

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

6

Countries and Territories with Active Zika Virus Transmission



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

7

Zika Virus in the USA

- Local transmission not yet reported in the continental US
- Zika is a nationally notifiable disease
- As of March 2, 153 travel-associated cases reported to CDC
- Nine pregnant travelers
 - 2 early pregnancy losses
 - 2 elective terminations
 - 1 infant with severe microcephaly
 - 2 healthy infants

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

8

Zika Virus in the USA

- With current outbreaks in the Americas, cases will increase
- Imported cases may lead to virus introduction and local spread in some areas of the US

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

9

Zika Virus Incidence and Attack Rates

- Yap Island outbreak, 2007 (population 7,391)
 - Infection rate: 73%
 - Symptomatic attack rate among infected: 18%
 - All age groups affected
 - Adults more likely to present for medical care
 - No severe disease, hospitalizations, or deaths



Duffy, M. N Engl J Med 2009

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

10

Zika Virus Vectors

- *Aedes* species mosquitoes.
 - *Aedes aegypti* – more efficient vector for humans.
 - *Aedes albopictus* – possible vector.
- Also transmit dengue and chikungunya viruses.
- Aggressive biters with peak feeding at daytime.
- Lay eggs in and around standing water.
- Live indoors and outdoors near households.
- Humans are the primary amplifying host during outbreaks.
- Monkeys are the natural reservoir

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

11

Zika Vectors





Aedes aegypti



Aedes albopictus

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

12



Other Modes of Transmission

Maternal-fetal: during pregnancy and time of birth.
Other documented modes of transmission: rare?

- Sexual
 - Male to female
 - One report of virus detected in semen at 62 days after onset
- Blood transfusion
 - Reports in Brazil being investigated
 - Deferral for 4 weeks in US
- Laboratory exposure

Theoretical concerns:

- Organ or tissue transplantation
 - Reports in Brazil being investigated
- Breast milk
 - Infectious virus has been detected

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

What We Don't Know

- How long Zika stays in semen?
- Do asymptomatic men have Zika in semen?
- Can asymptomatic men transmit Zika?
- Can a women transmit Zika to sex partners?
- Can Zika be transmitted through oral or anal sex?
- Does sexual transmission pose a different risk of birth defects than mosquito-borne transmission?

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

Clinical Disease Course

- Incubation several days to a week
- 80% asymptomatic
- Usually mild disease
 - Lasting several days to a week
- Hospitalization uncommon
- Fatalities rare
- Guillain-Barré syndrome reported following suspected Zika virus infection

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

Clinical Symptoms

Symptoms	N (n=31)	%
Macular or papular rash	28	90%
Subjective fever	20	65%
Arthralgia	20	65%
Conjunctivitis	17	55%
Myalgia	15	48%
Headache	14	45%
Retro-orbital pain	12	39%
Edema	6	19%
Vomiting	3	10%

Yap Island, 2007
Duffy M. N Engl J Med 2009

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

Need to Distinguish Zika from Dengue and Chikungunya

- All transmitted by the same mosquitoes with similar ecology
- Dengue and chikungunya can circulate in same area and rarely cause co-infections
- All have similar clinical features
- Important to rule out dengue, as proper clinical management can improve outcome

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

19

Treatment

- No specific antiviral treatment is available.
- Treatment is supportive: rest, fluids, and supportive care.
 - Assess for dengue and chikungunya viruses and avoid use of aspirin and other nonsteroidal anti-inflammatory drugs (NSAIDs) until dengue is ruled out (to reduce the risk of hemorrhage).
 - Treat pregnant women with acetaminophen.

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

20

Clinical Features: Zika Virus Compared to Dengue and Chikungunya

Features	Zika	Dengue	Chikungunya
Fever	++	+++	+++
Rash	+++	+	++
Conjunctivitis	++	-	-
Arthralgia	++	+	+++
Myalgia	+	++	+
Headache	+	++	++
Hemorrhage	-	++	-
Shock	-	+	-

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN


21

Zika and Microcephaly in Brazil

- Zika virus infections may be associated with microcephaly:
 - Considerable increase in the number of infants born with microcephaly has been observed in outbreak in Brazil during 2015.
 - True baseline rate of microcephaly and the association with Zika virus is unknown at this time.
 - Zika virus RNA isolated from several babies born with microcephaly and from fetal losses among women infected during pregnancy.
 - Some of the infants with microcephaly have tested negative for Zika virus.
- Investigations are ongoing.

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

22



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

23

Zika and Associated Birth Defects

- Microcephaly
- Brain atrophy
- Cerebral and intraocular calcifications
- Abnormal formed or absent brain structures
- Cataracts

Evidence not yet definitive

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

24



What We Don't Know

- Causal relation between Zika virus and microcephaly and other adverse pregnancy outcomes?
 - Full spectrum of impact in affected infants?
 - Impact of severity of maternal infection
 - Does asymptomatic pose a risk?
 - Timing of the infection during pregnancy effect on risk of fetal abnormalities?
 - Magnitude of the risk?
- WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN 26

Recent Study

- 88 women living in Rio de Janeiro who developed rash within previous 5 days
- Zika RT-PCR on blood and urine
 - 72 (82%) were positive
 - Range of time of infection was 5-35 weeks gestation
 - Fetal ultrasonography in 42 Zika-positive women
 - Abnormalities found in 12 (29%)

Brasil, P. et al NEJM March 10, 2016

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN 27

Recent Study (cont)

- Adverse findings
 - 2 fetal deaths at 36 and 38 weeks
 - 5 fetuses with in utero growth restriction with or without microcephaly
 - 7 with ventricular calcifications or other CNS lesions
 - 7 with abnormal amniotic fluid volume or cerebral or umbilical artery flow

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN 28

Laboratory Testing


- Processing of clinical specimens for Zika virus should be performed at a minimum of BSL2 precautions
- Perform a risk assessment to determine if higher levels of biocontainment are required
 - e.g. Suspicion of Chikungunya virus



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN 29

Diagnostic Testing

- Real-time PCR
- IgM Serology
- Immunohistochemical staining



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN 30

Reverse Transcriptase Real-Time PCR

- Two targets
 - Screening, broadly reactive target
 - Target specific for the Asian strain
- Waiting for a CDC Emergency Use Authorized triplex PCR
 - Zika, Dengue 1-4, and Chikungunya

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

31

Types of Specimens for Zika Virus PCR Testing

- Serum
- CSF
- Amniotic fluid (collected after 15 weeks gestation)
- Placental and umbilical cord tissues (fixed or frozen)
- Cord blood

WSLH will be testing only serum specimens at this time

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

32


IgM Capture ELISA

- CDC EUA assay
- Serum and CSF
 - CSF must be accompanied by a serum specimen
- IgM detectable ≥ 4 days after illness onset
 - Detectable up to 12 weeks

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

33

CDC IgM Capture ELISA



1. Coat With Goat anti-Human IgM
 - 4° Overnight
2. Add Patient Serum @ 1:400
 - 37° 1 Hour
3. Add Zika Antigen
 - 4° Overnight
4. Add HRP anti-Flavivirus McAb
 - 37° 1 Hour
5. Add substrate
 - RT 10 min
6. Add stop solution and Read

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

34

IgM Capture ELISA

- Difficult to distinguish infecting virus in people previously infected or vaccinated against a related flavivirus
- Anti-dengue virus IgM antibodies cross-react, so positive Zika IgM specimens must be confirmed
 - Plaque reduction neutralization assay (PRNT) performed at CDC

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

35

Diagnosis and Testing, Wisconsin

- All requests for fee-exempt Zika virus testing must be approved by the DPH. Phone 608-267-9003
- **Criteria for testing**
 - History of travel to an area with localized Zika virus transmission
 - Signs and symptoms within 2 weeks after returning. (Males or Females)
 - Asymptomatic pregnant females who are within 2-12 weeks of return from travel to an area with localized Zika virus transmission



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

36

Zika Testing Performed

History of Travel to area of Zika virus transmission AND Symptomatic

	PCR	IGM
Specimen collected within 3 days of onset	Zika Chik Dengue	
Specimen collected within 4-7 days of onset	Zika Chik Dengue	Zika Chik Dengue
Specimen collected within 1-12 weeks of onset		Zika Chik Dengue

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

37

Zika Testing Performed

Asymptomatic pregnant patient with history of travel to area of Zika virus transmission AND within 2-12 weeks of return from travel

Perform Only Zika virus IgM

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

38

Testing in Infants with Microcephaly or Intracranial Calcifications

- RT-PCR
 - Umbilical cord serum
 - Serum directly from infant within 2 days of birth
 - Maternal serum
 - Placental and cord tissue
 - CSF obtained for other studies
 - Mother's serum, saliva, and urine if not previously tested
- IgM ELISA
- IHC and histopathology on placenta and umbilical cord tissues

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

39

Testing in Infants without Microcephaly or Intracranial Calcifications

- If mother's test results positive or inconclusive
 - RT-PCR
 - IgM ELISA
 - IHC

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

40

Surveillance in Wisconsin

- Provide fee-exempt testing of Zika virus (also chikungunya and dengue) in travelers with appropriate signs and symptoms within 2 weeks after returning from areas with localized Zika virus transmission. Provide testing for asymptomatic pregnant women.
- Provide funding to assist the Wisconsin State Laboratory of Hygiene to bring on testing.
- Support mosquito surveillance for possible emerging *Aedes* species by collaborating with the University of Wisconsin-Madison, Medical Entomology Laboratory.
- Report all Zika virus confirmed and probable cases in real time to CDC ArboNet via WEDSS and National Electronic Disease Surveillance System (NEDSS).

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

41

Disease Reporting and Investigation

- Suspected Zika virus and other arboviral infections are Category II diseases and must be reported to public health within 72 hours: <https://www.dhs.wisconsin.gov/disease/diseasereporting.htm>
- Follow arboviral management protocol posted on the DHS website. <https://www.dhs.wisconsin.gov/publications/p0/p00894.pdf>
- Use Arboviral case report form for investigation.
- Disease reported category in WEDSS: Arboviral diseases, Zika virus.

Reminder: when investigation is completed for all arboviral diseases, the local health department should "send to state" for review, leaving the resolution status as "suspect". The DHS epidemiologist will determine if it meets the case definition for confirmed or probable case. Once the case is submitted as a confirmed or probable, it will be sent to CDC via NEDSS within 15-20 minutes.

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

42

Prevention

- Vaccines under development
- Avoid exposure to mosquitoes: use air conditioning or window/door screens; wear long sleeves and pants; use permethrin-treated clothing and gear, and Environmental Protection Agency (EPA)–registered repellents when outdoors.
- Pregnant women should consider postponing travel to any areas where Zika virus transmission is ongoing.
- Persons infected with Zika, dengue, or chikungunya viruses should be protected from further exposure to mosquitoes during illness to reduce the risk of local transmission.

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

43

Prevention of Sexual Transmission

- Men who reside in or have traveled to an area of active Zika virus transmission who have a pregnant partner **should**
 - Abstain from sexual activity or use condoms during sex
- Men with nonpregnant sex partners **might consider** abstaining or use condoms

MMWR/February 12, 2016/Vol. 65/No. 5

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

44

Wisconsin Residents Tested

As of March 8, 2016

	Patients	Percentage
Asymptomatic	159	73%
Symptomatic	58	27%
Total	217	

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

45


Research at UW Madison

- Drs. Dave O'Connor and Jorge Osorio
 - <https://dholk.primat.wisc.edu/project/dho/public/Zika/public/begin.view>
- Infected 3 rhesus monkeys subcutaneously with 10^6 , 10^5 , and 10^4 PFU of Zika virus
- Looking out
 - Viral RNA quantification
 - Blood counts
 - Immunology
 - Blood Chemistry

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

46


Plasma Viral Loads



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

47


Urine Viral Loads



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

48

CSF Viral Loads



WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

49

References and Resources

- CDC COCA. Zika Virus — What Clinicians Need to Know: http://emergency.cdc.gov/coca/calls/2016/callinfo_012616.asp
- CDC MMWR. Zika Virus Spreads to New Areas — Region of the Americas, May 2015–January 2016: <http://www.cdc.gov/mmwr/volumes/65/wr/mm6503e1.htm>
- CDC MMWR. Interim Guidelines for Pregnant Women During a Zika Virus Outbreak — United States, 2016: <http://www.cdc.gov/mmwr/volumes/65/wr/mm6502e1.htm>
- Wisconsin Division of Public Health – Zika Virus Webpage: <https://www.dhs.wisconsin.gov/arboviral/zika.htm>
- CDC Zika Virus Webpage: <http://www.cdc.gov/zika/index.html>

WISCONSIN STATE LABORATORY OF HYGIENE - UNIVERSITY OF WISCONSIN

50